SCREENING QUESTIONNAIRE

Binocular Vision Dysfunction Questionnaire (BVDQ™)



Directions: For each of the following questions, please check the answer that best describes your situation. If you wear glasses or contact lenses, answer the questions assuming that you are wearing them.

	Always = every day Occasionally = less than once per week	Frequently = at least once per week Never = never	W. Aller		MO, STAN
1	Do you have headaches and/or facial pain?				
2	Do you have pain in your eyes with eye movement?				
3	Do vou experience neck or shoulder discomfort?				

	bo you have neadaches and of facial pain.							
2	Do you have pain in your eyes with eye movement?							
3	Do you experience neck or shoulder discomfort?							
4	Do you have dizziness and/or light headedness?							
5	Do you experience dizziness, light headedness, or nausea while performing close-up activities (computer work, reading, writing, etc.)?							
6	Do you experience dizziness, light headedness or nausea while performing far-distance activities (driving, television, movies, etc.)?							
7	Do you experience dizziness, light headedness, or nausea when bending down and standing back up, or when getting up quickly from a seated position?							
8	Do you feel unsteady or drift to one side while walking?							
9	Do you feel overwhelmed or anxious while walking in a large department store (Target, Wal-Mart, Costco, etc.)?							
10	Do you feel overwhelmed or anxious when in a crowd?							
11	Does riding in a car make you feel dizzy or uncomfortable?							
12	Do you experience anxiety or nervousness because of your dizziness?							
13	Do you ever find yourself with your head tilted to one side?							
14	Do you experience poor depth perception or have difficulty estimating distances accurately?							
15	Do you experience double/overlapping/shadowed vision at far distances?							
16	Do you experience double/overlapping/shadowed vision at near distances?							
17	Do you experience glare or have sensitivity to bright lights?							
18	Do you close or cover one eye with near or far tasks?							
19	Do you skip lines or lose your place when you are reading? Do you use your finger, ruler or other guides to maintain your position on the page?							
20	Do you tire easily with close-up tasks (computer work, reading, writing)?							
21	Do you experience blurred vision with far-distance activities (driving, television, movies, chalkboard at school, etc.)?				Г			
22	Do you experience blurred vision with close-up activities (computer work, reading, writing, etc.)?							
23	Do you blink to 'clear up' distant objects after working at a desk or working with close-up activities (computer work, reading, writing, etc.)?							
24	Do you experience words running together while reading?							
25	Do you experience difficulty with reading or reading comprehension?							

For questions 1 - 25 on the front, scoring is as follows (see below). Add the scores for questions 1 - 25 to get a TOTAL score.							
Always = $ x 3 $	Frequently =	_x2	Occasionally =	x1	Never=	x0	TOTAL (BVDQ)
		A BV	DQ score of 15 or gre	ater is	highly sugges	stive of a	Binocular Vision Dysfunction.

		None	Worst		None	Worst
On an average day, how	Dizziness	0 1 2 3 4 5 6	7 8 9 10	Neckache	0 1 2 3 4 5 6 7 8	3 9 10
much are you bothered by symptoms listed here?	Nausea	0 1 2 3 4 5 6	7 8 9 10	Unsteady when walking	0 1 2 3 4 5 6 7 8	3 9 10
Rate each symptom from $0-10$ 0 = None of that symptom	Anxiety	0 1 2 3 4 5 6	7 8 9 10	Sensitivity to light	0 1 2 3 4 5 6 7 8	3 9 10
10 = Worst	Headache	0 1 2 3 4 5 6	7 8 9 10	Reading difficulty	0 1 2 3 4 5 6 7 8	3 9 10
				Sound sensitivity	0 1 2 3 4 5 6 7 8	3 9 10

For the above questions, total up the 9 numbers above and total them to the right. This score is called a **SSI** (or **Symptom Severity Index**). An SSI Score of **15 or greater** is highly suggestive of a Binocular Vision Dysfunction.

TOTAL (SSI)

Name	Date
Phone Number	Email
Please record any additional symptoms you may be experiencing or spe	ecific concerns that you have about your eyes/vision:

This questionnaire is designed to screen for those who may have difficulty with vision alignment. The information obtained herein is considered a preliminary result only and does not diagnose or constitute confirmation of any vision problems. It is not a substitute for a NeuroVisual examination. Since vision changes can occur without visible indications, most eye care professionals and medical authorities recommend a vision exam annually